arXiv:cond-mat/9612017v1 [cond-mat.mes-hall] 2 Dec 1996

Thermal activation of exclusonic quasiparticles in the fractional quantum Hall effect

J. Shiraishi[1] , M. Kohmoto[1] and Y.S. Wu[2]
[1]*Institute for Solid State Physics, University of Tokyo, Roppongi, Minato-ku, Tokyo 106, Japan*
[2]*Department of Physics, University of Utah, Salt Lake City, UT 84112, U.S.A.*

Abstract

Quasiparticles of the fractional quantum Hall systems obey fractional (including mutual) exclusion statistics. In this note we study the effects of exclusion statistics on thermal activation of quasiparticle pairs in the approximation of generalized ideal gas. The distribution function for thermally activated quasiparticles is found explicitly for the statistics matrix given by the composite fermion picture and the thermodynamic consequences are discussed analytically. In particular, at low temperatures, there is a quasielectron-quasihole duality in the description of the system between two adjacent magic fillings $\nu(m,p) \equiv m/(2pm+1)$ and $\nu(m+1,p)$.

Typeset using REVTEX

A fascinating aspect of the fractional quantum Hall (FQH) effect is that quasiparticles in the system have exotic quantum numbers, very different from those of the constituent particles, the electrons. For example, the gapful FQH liquid with electron filling close to $\nu = m/(2pm + 1)$ (p and m being integers) has two species of quasiparticle excitations (quasielectrons labeled by $+$ and quasiholes labeled by $-$) [1], both of which are fractionally charged ($e^*_\pm = \mp 1/(2pm + 1)$) anyons [2,3] with fractional exchange statistics [4]. Furthermore, they satisfy novel quantum statistics in the sense of state counting, which is neither Bose or Fermi. In this note we study the effects of quantum statistics on thermal activation of the FQH quasiparticle pairs and their thermodynamic consequences.

Previously it was known that a single state can not be occupied by more than one fermions; on the other hand, any number of bosons can occupy a single state. Haldane [5] first proposed that in some strongly correlated many-body systems, the system may be described by quasiparticles that obey *fractional exclusion statistics* other than Bose or Fermi statistics. We call such particles or quasiparticles as "*exclusons*". One of us [6] has considered generalized ideal gas of non-interacting exclusons, and formulated their statistical distribution and thermodynamics. (Note that this does not mean the interactions are neglected. For example, there are one-dimensional exactly solvable models which can be precisely represented as a gas of non-interacting exclusons; see *e.g.* [8–10]). In refs. [5,11–14] it is shown that quasiparticles in the FQH systems with electron filling $\nu = 1/(2p+1)$ obey fractional exclusion statistics, including mutual exclusion statistics between quasielectrons and quasiholes. In ref. [6] the low-temperature thermodynamics of FQH quasiparticles in these systems is discussed in the framework of generalized ideal gas. In this note we consider the more general FQH systems close to electron filling $\nu(m,p) \equiv m/(2pm + 1)$. One way to study the effects of exotic exclusion statistics, especially the effects of mutual statistical exclusion between quasielectrons and quasiholes, is to see how they affect thermal activation of quasiparticle pairs. When the statistics matrix of quasiparticles is the one determined by the composite fermion picture [16], we have been able to find explicit distribution functions for quasiparticles at finite temperature and to discuss analytically the resulting thermodynamics. Such an analytic solutions have not been available even for one-dimensional solvable models. In particular, we find a quasiparticle-quasihole duality in the quasiparticle gas at low temperatures: A system with ν between two adjacent magic fillings $\nu(m,p)$ and $\nu(m+1,p)$ can be equivalently described either in terms of quasielectrons in the FQH fluid with $\nu(m,p)$ or in terms of quasiholes in the FQH fluid with $\nu(m+1,p)$.

In Jain's composite fermion approach, the fractional quantum Hall (FQH) state of electrons in physical magnetic field is explained as the integer quantum Hall (IQH) state of composite fermions in an effective magnetic field [16]. Imagine an adiabatic process, by which somehow a certain amount of magnetic flux is collected to electrons, so that finally $2p$ flux quanta (p a positive even integer) are attached to each electron to form a composite fermion. These composite fermions are now moving in a reduced effective magnetic field $B_{eff} = B - 4\pi p\rho$, where ρ is the number density of electrons (or composite fermions). (In our convention, the unit of flux quantum is 2π.) Therefore the effective filling factor for composite fermion has been increased to ν_{eff}, given by $\nu_{eff}^{-1} = (B - 4\pi p\rho)/2\pi\rho = \nu^{-1} - 2p$. For $\nu_{eff} = m$ (with m an integer), we have $\nu^{-1} = 2p + m^{-1}$ and $\nu = m/(2pm + 1)$. Thus, fractional quantum Hall systems with $\nu(m,p) \equiv m/(2pm+1)$ are adiabatically changed into an integer Hall system with filling factor m, as was emphasized by Greiter and Wilczek [17].

Let us consider quasiparticles, quasielectrons and quasiholes, in the FQH liquid with filling close to $\nu_0 = m/(2pm+1)$. Let $N_\phi \equiv eBV/hc$, N_+ and N_- be the number of total flux quanta, the number of quasielectrons and quasiholes, respectively. Since quasielectrons and quasiholes are fractionally charged ($e^*_\pm = \mp 1/|2pm+1|$), the total charge balance means $N_e = \nu_0 N_\phi + |e^*_+|N_+ - |e^*_-|N_-$, and the filling ν is given by

$$\begin{aligned}\nu \equiv \frac{N_e}{N_\phi} &= \nu_0 + |e^*_+|\frac{N_+}{N_\phi} - |e^*_-|\frac{N_-}{N_\phi} \\ &= \frac{m}{2pm+1} + \frac{1}{(2pm+1)^2}(n_+ - n_-)\ ,\end{aligned} \tag{1}$$

where $n_\pm$ is the average "occupation number per state" $n_\pm = N_\pm/|e^*_\pm|N_\phi = |2pm+1|N_\pm/N_\phi$, which is related to the density of quasiparticles $\rho_\pm(T)$ by $\rho_\pm(T) = |e^*_\pm|N_\phi/Vn_\pm$. In what follows, we restrict the range of the filling factor as $\nu < 1/2p$ (or $\nu > 1/2p$) if $m \geq 0$ (or $m \leq -1$).

Let us count states of quasiparticles in the FQH system with filling near $\nu_0 = m/(2pm+1)$. The effective flux for the composite fermions is given by $N_{\phi,eff} = N_\phi - 2pN_e$. At ν_0 the ground state of the system corresponds to $|m|$ filled Landau levels for composite fermions. Note that $\mathrm{sgn}(N_{\phi,eff}) = \mathrm{sgn}(m)$. We will consider the temperature range that is low enough so that the quasiparticle excitations of the system are dominated by composite fermions excited into their $(|m|+1)$-th Landau level. Thus, quasielectrons correspond to composite fermions in the $(|m|+1)$-th Landau level, while quasiholes holes of composite fermions in the filled $|m|$-th Landau level. If the composite-fermion excitations, which should be considered as unit-charged for consistency, obey usual Fermi statistics, then the number of available single-particle states for them are obviously

$$G^{(b)}_{eff,\pm} = |N_{\phi,eff}| - (N_\pm - 1). \tag{2}$$

To derive the exclusion statistics of the quasiparticle excitations, we need to express $N_{\phi,eff}$ in (2) in terms of the external N_ϕ, resulting in

$$\begin{aligned}G^{(b)}_{eff,+} &= G_+ - \alpha_{++}N_+ - \alpha_{+-}N_-, \\ G^{(b)}_{eff,-} &= G_- - \alpha_{-+}N_+ - \alpha_{--}N_-,\end{aligned} \tag{3}$$

where $G_\pm = |e^*_\pm|N_\phi$ is the single-quasiparticle degeneracy in terms of the external flux with the proportionality constant $1/|2pm+1|$ identified as the fractional charge (absolute value) of the quasiparticles, and the exclusion statistics matrix $\alpha_{\pm\pm}$ for quasiparticles are:

$$\alpha_{\epsilon\epsilon'} = \delta_{\epsilon,\epsilon'} + \epsilon' \frac{2p}{2pm+1}. \tag{4}$$

We emphasize that these values of statistics are derived under the assumption of eq. (2), which looks very natural in the composite fermion picture. (However, modification is not absolutely impossible. We leave this problem to discussions at the end of the paper.)

The diagonal statistics satisfies $\alpha_{--}(m+1) = 1/\alpha_{++}(m)$. This "symmetry" becomes important while we study the duality in the quasiparticle gas at low temperatures.

When the temperature is low, the density of thermally activated quasiparticle pairs is not very high, so that quasiparticle interactions other than statistical ones may be ignored in the first approximation. The system is therefore described by a generalized ideal gas [6] with two species, for which the following two conditions are satisfied:
1) The total energy (eigenvalue) is always of the form of a simple sum, in which the i-th term is linear in the particle number N_i: $E = \sum_i N_i \varepsilon_i$,
with ε_i identified as the energy of a quasielectron ($i = +$) or a quasihole ($i = -$).
2) The following state-counting for statistical weight W, i.e. the dimension of the Hilbert space of quasiparticle states, is applicable:

$$W = \begin{pmatrix} G^b_{eff,+} + N_+ - 1 \\ N_+ \end{pmatrix} \begin{pmatrix} G^b_{eff,-} + N_- - 1 \\ N_- \end{pmatrix} . \tag{5}$$

Consider a grand canonical ensemble at temperature T and with chemical potential $\mu_i (i = +, -)$ for species i, whose partition function is given by (with k the Boltzmann constant)

$$Z = \sum_{\{N_i\}} W(\{N_i\}) \ \exp\{\sum_i N_i(\mu_i - \varepsilon_i)/kT\} \ . \tag{6}$$

where W is the statistical weight (5). One obtains the equations determining the most probable distribution of $n_\pm$:

$$\sum_j \alpha_{ij} n_j(T) = 1 - w_i(T) \ , \tag{7}$$

with $w_\pm(T)$ being determined by the functional equations

$$\begin{aligned} {w_+}^{\alpha_{++}}(1+w_+)^{1-\alpha_{++}}\Big(\frac{w_-}{1+w_-}\Big)^{\alpha_{-+}} &= e^{(\varepsilon_+ - \mu_+)/kT} \ , \\ {w_-}^{\alpha_{--}}(1+w_-)^{1-\alpha_{--}}\Big(\frac{w_+}{1+w_+}\Big)^{\alpha_{+-}} &= e^{(\varepsilon_- - \mu_-)/kT} \ . \end{aligned} \tag{8}$$

By solving (7), $n_\pm$ are expressed in terms of $w_\pm$ as

$$\begin{aligned} n_+(T) &= \frac{w_- + \alpha_{--} - \alpha_{+-}}{(w_+ + \alpha_{++})(w_- + \alpha_{--}) - \alpha_{+-}\alpha_{-+}} \ , \\ n_-(T) &= \frac{w_+ + \alpha_{++} - \alpha_{-+}}{(w_+ + \alpha_{++})(w_- + \alpha_{--}) - \alpha_{+-}\alpha_{-+}} \ . \end{aligned} \tag{9}$$

These equations can be solved explicitly as

$$\begin{aligned} w_+(T) = \frac{1}{2f(\nu, m-1)}\Big(& -(e^{\Delta/kT}+1) f(\nu, m) \\ & +\mathrm{sgn}(m)\sqrt{(e^{\Delta/kT}+1)^2 f(\nu,m)^2 - 4e^{\Delta/kT} f(\nu, m+1) f(\nu, m-1)}\Big), \end{aligned} \tag{10}$$

$$\begin{aligned} w_-(T) = \frac{1}{2f(\nu, m+1)}\Big(& -(e^{\Delta/kT}+1) f(\nu, m) \\ & -\mathrm{sgn}(m)\sqrt{(e^{\Delta/kT}+1)^2 f(\nu,m)^2 - 4e^{\Delta/kT} f(\nu, m+1) f(\nu, m-1)}\Big), \end{aligned} \tag{11}$$

where $f(\nu, m) \equiv (2pm+1)\nu - m$, $\Delta = \varepsilon_+ + \varepsilon_-$ and we have used the charge conservation $\mu_+ + \mu_- = 0$. The sign functions in front of the square roots are suitably chosen so that we have positive $w_i(T)$ for $\nu \sim \nu_0$. It is remarkable that a closed expression for $w_i(T)$ is available for our model of FQH systems. This is closely related to the structure of the statistics matrix given by eq. (4). In Fig.1, $N_\pm(T)$ are plotted against ν. We have quite remarkably simple form: $n_\pm(T)|_{\nu=\nu_0} = 1/(1+e^{\Delta/2kT})$, thus we have

$$\rho_\pm(T)\Big|_{\nu=\nu_0} = \frac{|e^*_\pm| N_\phi}{V} \frac{1}{1+e^{\Delta/2kT}}. \tag{12}$$

It may be possible to check experimentally whether our results and the fractional charges are correct or not.

Now we proceed to consider thermodynamic consequences of the above analytic distribution function. First, the entropy is given by

$$S = \frac{N_\phi}{k|2pm+1|} \sum_{i=\pm} \Bigg((1 + n_i - \sum_{j=\pm} \alpha_{ij} n_j)\ \log(1 + n_i - \sum_{j=\pm} \alpha_{ij} n_j) \\ - n_i \log n_i - (1 - \sum_{j=\pm} \alpha_{ij} n_j)\ \log(1 - \sum_{j=\pm} \alpha_{ij} n_j) \Bigg). \tag{13}$$

From this expression, we note a duality of the system which becomes exact in the zero temperature limit. From now on, we assume $m > 0$ and fix p for simplicity .

Let us study the entropy for the system with filling in the region $m/(2pm+1) \leq \nu \leq (m+1)/[2p(m+1)+1]$ in two ways. It can be shown that at low temperatures, the approximate equality

$$S(m,p) \sim S(m+1,p) \tag{14}$$

holds. This indicates that at low temperatures, the same system with $m/(2pm+1) \leq \nu \leq (m+1)/[2p(m+1)+1]$ can be equivalently described either in terms of quasielectrons in the FQH fluid with $\nu(m,p)$ or in terms of quasiholes in the FQH fluid with $\nu(m+1,p)$ See Fig. Note that at zero temperature we have an exact duality $S(m,p) = S(m+1,p)$.

We also note that in between two adjacent magic fillings $\nu(m,p)$ and $\nu(m+1,p)$, there is a filling at which the entropy has a local maximum. This filling can be obtained by solving $\partial_\nu S(m,p)|_{T\to 0} = 0$. In Table 1, we list some of the fillings where the entropy has local maximum. Physically, it is believed that the transition between plateaus of different quantized Hall conductance σ_{xy} is due to quantum percolation transition [18] of quasielectrons (or quasiholes from the dual point of view). It is natural to expect that the entropy should be a maximum at the percolation transition point, so one may interpret the solution of $\partial_\nu S|_{T\to 0} = 0$. as the approximate position of the transition point, if the percolation transition persists in the low disorder limit (see Table 1).

Second, the thermodynamic potential $\Omega = -kT \log Z$ is [6]

$$\Omega \equiv -PV = -kT \sum_i G_i \log(1 + \frac{n_i}{1 - \sum_j \alpha_{ij} n_j}) \\ = -\frac{kT N_\phi}{|2pm+1|} \log \frac{1+w_+}{w_+} \frac{1+w_-}{w_-} \tag{15}$$

Using the analytic expressions (10) and (11), we have the following equation of state:

$$PV = \frac{kTN_\phi}{|2pm+1|}\log\left\{\frac{(2p\nu-1)}{e^{\Delta/kT}f(\nu,m+1)f(\nu,m-1)}\left(-(1+e^{\Delta/kT})(2p\nu-1)\right.\right. \tag{16}$$
$$\left.\left.+\mathrm{sgn}(m)\sqrt{(1+e^{\Delta/kT})^2f(\nu,m)^2-4e^{\Delta/kT}f(\nu,m+1)f(\nu,m-1)}\right)\right\}.$$

Examining this expression for the pressure, we can see how new incompressible FQH fluid states arise in thermodynamics: They emerge at fillings where the pressure becomes divergent (see Fig. 3). It can be shown that $P>0$ in the region $\nu_0(m-1,p)\le\nu\le\nu_0(m+1,p)$ and at finite temperatures P diverges at temperature-independent fillings $\nu=\nu_0(m\pm1,p)$. The divergence of the pressure, of course, is due to the complete filling (of composite fermions) of a new Landau level, or the quasiparticle's condensation into a new incompressible FQH state, at $\nu_0^{\rm new}=(m\pm1)/(2p(m\pm1)+1)$.

Note that $P(m,p)\ne P(m+1,p)$ even at low temperature. Since the pressures $P(m,p)$ and $P(m+1,p)$ are measured from different incompressible ground states, this is not contradictory to the duality of the system.

To conclude, we make the following remarks. If the filling factor approaches the value with the even denominator $(m\to\infty)$ $\nu=1/2p$, the statistics matrix degenerates to that of free fermion. So, at this filling the quasiparticles are nothing but free fermions. However, the thermodynamics for the $\nu=1/2p$ state may not be that simple. Since if we allow fluctuations of the filling fraction near $1/2p$, as in a grand canonical ensemble, we have to do averaging over many FQH states around the filling $1/2p$. It seems important to study this averaging effect.

Also, we note that our results crucially depend on the assumption (2). Though this seems very natural in the (unprojected) composite fermion picture, there is numerical data for small systems on a sphere [14,15], whose interpretation requires modification of eq. (2) to

$$G^{(b)}_{eff,\pm}=|N_{\phi,eff}|-(N_\pm-1)\pm N_\mp. \tag{17}$$

This will change eq. (4) to $\alpha_{++}=2-\alpha_{--}=\alpha_{-+}=-\alpha_{+-}=1+2p/(2pm+1)$. Despite the numerical evidence in favor of this modified parameters, it may be possible , for example using (12), to choose between these from experimental data.

ACKNOWLEDGMENTS

It is a pleasure to thank S. Katsumoto for useful discussions. The work of Y.S.W. was supported in part by U.S. NSF grant PHY-9601277.

REFERENCES

[1] R.B. Laughlin, *Phys. Rev. Lett.* **50**, (1983) 1359.

[2] B.I. Halperin, *Phys. Rev. Lett.* **52** , (1984) 1583.

[3] D. Arovas, J.R. Schrieffer and F. Wilczek, *Phys. Rev. Lett.* **53**, (1984) 722.

[4] See, e.g., papers collected in "*Fractional Statistics and Anyon Superconductivity*", ed. F. Wilczek, World Scientific (Singapore), 1989; and references therein.

[5] F.D.M. Haldane, *Phys. Rev. Lett.* **67**, (1991) 937.

[6] Y.S. Wu, *Phys. Rev. Lett.* **73**, (1994) 922.

[7] S.B. Isakov, *Phys. Rev. Lett.* **73**, (1994) 2150.

[8] D. Bernard and Y.-S. Wu, in "*Yangians and Recent Developments on Quantum Integrable Models*", Proceedings of the Sixth Nankai Workshop (June 1994, Tianjin, China), edited by M.L. Ge and Y.S. Wu; World Scientific, Singapore, 1995.

[9] S.B. Isakov, *Int. J. Mod. Phys.* **A9**, (1994) 2563.

[10] Y. Hatsugai, M. Kohmoto, T. Koma and Y.S. Wu, to be published in Phys. Rev. B.

[11] S. He, X.C. Xie and F.C. Zhang, *Phys. Rev. Lett.* **68**, (1992) 3460.

[12] J. Yang and W.P. Su, *Phys. Rev. Lett.* **70**, (1993) 1163.

[13] G.S. Canright and M.D. Johnson, *Phys. Rev.* **B 49**, (1994) 2947.

[14] W.P. Su, Y.S. Wu and J. Yang, *Phys. Rev. Lett.* **77**, 3423.

[15] S.B. Isakov, G.S. Canright and M.D. Johnson, preprint cond-mat/9608139.

[16] J.K. Jain, *Phys. Rev. Lett.* **63**, (1989) 199.

[17] M. Greiter and F. Wilczek, *Mod. Phys. Lett.* **4**, 1063 (1990).

[18] D.-H. Lee, Z. Wang and S. Kivelson, *Phys. Rev. Let.* **70** , (1993) 4130.

FIGURES

1 Quasiparticle densities $n_{\pm}(T)$ with $m = 2, p = 1$ are plotted for $1/3 < \nu < 3/7$ at the temperature $e^{\Delta/kT} = 100$.

2 Entropy corresponding to two neighboring parameters are plotted: $S(m = 2)k/N_{\phi}$ and $S(m = 3)k/N_{\phi}$ for $1/3 < \nu < 4/9$ at the temperature $e^{\Delta/kT} = 100$. It is observed that $S(m = 2) \sim S(m = 3)$ in the region $2/5 < \nu < 3/7$.

3 Plot of PV/kTN_{ϕ} for $1/3 < \nu < 3/7$ at the temperature $e^{\Delta/kT} = 100$. Note the divergence of the pressure at $\nu = 1/3$ and $3/7$.

TABLES

models with $m \geq 1$, $p = 1$	$\nu = 0.3692$	0.4150	0.4368	0.4497
$m \leq -1$, $p = 1$	0.8057	0.6307	0.5850	0.5632
$m \geq 1$, $p = 2$	0.2121	0.2267	0.2331	0.2367
$m \leq -1$, $p = 2$	0.3064	0.2788	0.2696	0.2648

TABLE I. Some of the transition points obtained from the condition $\partial_\nu S|_{T \to 0} = 0$ are listed.